



1/7

FIG. 1



2/7

FIG. 2 TOP RIGHT SLEEVE UNDONE
TO ENABLE ARM WITHDRAWAL



3/7

FIG. 4

ARM PASSED THROUGH
SLEEVE PRIOR TO
ACCESS TO RIGHT CHEST





FIG. 3


ANAESTHETIZED PATIENT
ARM BEGINNING TO BE
WITHDRAWN FROM SLEEVE
WITHOUT LIFTING PATIENT



4/7


FIG. 5

ANAESTHETIZED PATIENT ARM FULLY
WITHDRAWN FROM SLEEVE WITHOUT
LIFTING PATIENT
CHEST EXPOSED FOR ANAESTHETIST




5/7

FIG. 6



6/7



SUGISUIT SHORTS UNDONE TO
ALLOW ACCESS TO RIGHT LEG
AND GROIN, VELCRO UNDONE

FIG. 7A




FIG. 7B





FIG. 7C

7/7



VELCRO UNDONE BOTH LEGS
FRONT FLAP LIFTED DOWN
ACCESS TO RIGHT AND LEFT LEG
WITHOUT LIFTING PATIENT
ELASTICATED TOP
VELCRO

FIG. 8A




FIG. 8B




FIG. 8C